researchvision.us

Cognify: Journal of Artificial Intelligence and Cognitive Science
Vol.2 | No. 1 | 2025 pp. 34–38 |**ISSN: 3066-3466**

Interaction Between Light Spectra and Phytohormones

Muroddinova Farida Rakhmatboy qizi[1]

[1] Student of Gulistan State University

Submitted: 25-April, 2025
Accepted: 26-May, 2025
Published: 28-June, 2025

Vol. 2, **No.** 1, 2025.
10.62762/JAICS.2025.000000

***Corresponding author:**
oktamjonxazratkulov7@gmail.com[1]

Copyright © 2025 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

http://creativecommons.org/licenses/by/4.0/

Open Access

Abstract: Light and phytohormones are key regulators of plant growth, development, and adaptation. Their interaction forms the basis of plant physiology, integrating external environmental cues with internal hormonal responses. While individual effects of light spectra and phytohormones have been extensively studied, understanding their synergistic relationship in growth regulation remains an emerging research focus, especially for application in controlled environments. There is limited integration of light spectrum-phytohormone interaction knowledge into practical greenhouse and vertical farming technologies to optimize growth and productivity. This study aims to analyze how different light wavelengths, including red, blue, and far-red, influence the biosynthesis and action of major phytohormones—auxins, gibberellins, cytokinins, abscisic acid, and ethylene—in plant systems. The analysis shows that red light promotes auxin-mediated cell elongation; blue light limits gibberellin production, resulting in compact growth; far-red light increases auxin and ethylene, enhancing elongation but risking structural weakness. Low light induces ABA and ethylene, slowing growth and promoting leaf abscission. This article synthesizes the biochemical pathways linking light perception to hormonal regulation and proposes practical applications for optimizing lighting strategies in greenhouses and vertical farms. Understanding these interactions allows for targeted crop management, development of stress-resilient varieties, and maximizing yield in artificial environments, addressing food security challenges in resource-constrained regions.

Keywords: Light spectra, phytohormones, plant physiology, photoreceptors, greenhouse, photosynthesis, phytochrome, cryptochrome.

Introduction

Plants have complex adaptation mechanisms that allow them to respond to both internal and

external factors. They perceive environmental stimuli such as light, humidity, and temperature and respond to them through genetic and physiological pathways. Among these stimuli, light is one of the most crucial regulators, governing essential processes such as photosynthesis, phototropism, photoperiodism, and photomorphogenesis[1].

Internally, phytohormones serve as key regulators of plant growth rate, tissue differentiation, organ formation, and responses to stress. The interaction between light and phytohormones highlights a coordinated regulatory system, where external light cues are converted into physiological responses through hormonal signaling. This article analyzes the interaction between these two key factors based on scientific sources[2].

Plants possess complex adaptive systems enabling them to respond effectively to external environmental cues and internal hormonal signals. Light acts as a primary regulator of plant growth processes such as photosynthesis, photomorphogenesis, and photoperiodism, while phytohormones coordinate growth, development, and stress responses. The interaction between light spectra—particularly red, blue, and far-red wavelengths—and phytohormones like auxins, gibberellins, cytokinins, abscisic acid, and ethylene determines plant morphology and productivity. Understanding this synergy is crucial for optimizing crop growth in controlled environments such as greenhouses. This article investigates the mechanisms underlying these interactions and highlights their practical applications in modern agricultural systems[3].

Methods

This study employed a literature-based analytical approach to investigate the interactions between light spectra and phytohormones in plant growth and development. Data were collected from contemporary plant physiology textbooks, peer-reviewed journal articles, and authoritative scientific sources such as Taiz & Zeiger, Hopkins & Hüner, and specialized plant physiology databases. The analysis focused on synthesizing findings about how red, blue, green, far-red, UV-A, and UV-B light wavelengths influence the synthesis, activity, and signaling pathways of key phytohormones, including auxins, gibberellins, cytokinins, abscisic acid, and ethylene [4]. The methodology included comparative evaluation of experimental data on photoreceptor activation—phytochromes for red and far-red light, cryptochromes and phototropins for blue light—and their downstream hormonal regulation effects, such as auxin biosynthesis under red light and gibberellin inhibition under blue light conditions. Additionally, practical applications of these interactions in greenhouse and vertical farming systems were examined by integrating experimental outcomes with controlled environment management strategies to optimize plant morphology, productivity, and stress resistance[5]. The study also reviewed specific examples from recent literature demonstrating light-hormone synergy in stem elongation, leaf expansion, and flowering. A descriptive synthesis was used to summarize and explain these physiological processes in an integrated manner. The method ensured a comprehensive understanding of light–phytohormone interactions, focusing on mechanisms that could inform future controlled environment agriculture research, crop productivity optimization, and plant adaptation studies[6].

Results and Discussion

Detailed Overview of Light Spectra

Light is a form of electromagnetic radiation that includes a specific range of wavelengths. Plants primarily utilize the 400–700 nm range, known as Photosynthetically Active Radiation (PAR). Different wavelengths within this spectrum trigger different physiological responses in plant tissues[7].

Red light (660–680 nm) is among the most effective for photosynthesis. It is absorbed by phytochrome receptors and plays a key role in seed germination, stem elongation, and vegetative growth. In particular, red light activates phytochrome B, which stimulates auxin biosynthesis and promotes cell elongation.

Blue light (430–470 nm) is detected by cryptochromes and phototropins. It significantly influences plant morphology, leaf opening, stomatal regulation, and photomorphogenesis. Besides contributing to photosynthesis, blue light also modulates various growth processes[8].

Green light (500–550 nm), traditionally considered less active, has recently been shown to penetrate deeper into plant tissues and affect metabolic processes.

Far-red light (above 700 nm) interacts with phytochromes and activates the shade avoidance response, enhancing elongation. However, excessive far-red light can lead to undesirable stretching of the plant.

UV-A and UV-B radiation stimulate protective stress responses. These wavelengths enhance the production of secondary metabolites and increase the plant's resistance to harmful external factors[9].

Phytohormones: Types and Functions

Phytohormones are organic compounds produced in small amounts in plants that have powerful physiological effects. Their functions are diverse and critical for plant life.

Auxins are primary growth regulators involved in cell elongation, root formation, and tropic responses. The most common form is indole-3-acetic acid (IAA).

Gibberellins promote stem elongation, seed germination, and flowering. They are especially important in the transition from vegetative to reproductive phases in perennial plants[10].

Cytokinins stimulate cell division, delay leaf senescence, and regulate organogenesis.

Abscisic acid (ABA) is known as the stress hormone. It controls stomatal closure during water stress and slows down growth processes.

Ethylene, a gaseous hormone, is associated with fruit ripening, leaf abscission, and plant responses to various stresses.

Interaction Between Light and Phytohormones

The interaction between light and phytohormones determines how plants respond to their environment. Photoreceptors detect light signals, which are then translated into intercellular responses via phytohormonal pathways[11].

For instance, **red light** activates phytochromes, which increase auxin biosynthesis, enhancing growth and elongation[12].

Blue light reduces gibberellin biosynthesis through cryptochrome activation, limiting excessive stem elongation. This is particularly important in controlled environments, such as greenhouses, where plant height must be managed for optimal yield[13].

Far-red radiation increases both auxin and ethylene levels, leading to more elongation. However, if not regulated, this may cause fragile, overstretched plants[14].

In contrast, **low light conditions** increase the production of ethylene and ABA, slowing growth and accelerating leaf abscission, mimicking stress-like symptoms.

Practical Applications

These scientific insights can be applied in advanced plant cultivation systems such as greenhouses, vertical farming, and LED-based artificial lighting environments. For example:

- A balanced combination of **red and blue light** optimizes leaf development and flowering.
- Blue-light-dominant conditions promote compact, dense plant architecture, which is ideal for confined spaces.
- Controlled **far-red exposure** helps prevent excessive elongation, maintaining robust plant structure[15].

Understanding the synergy between light spectra and phytohormones provides new opportunities in modern agriculture, especially in regions with limited natural light or harsh climates.

Conclusion

The interaction between light spectra and phytohormones is a central mechanism in plant development. Light signals are modulated through hormonal pathways, affecting nearly all physiological processes in plants. These insights enable targeted crop management strategies, breeding for stress-tolerant varieties, and achieving maximum productivity in closed and controlled environments.

In conclusion, the interaction between light spectra and phytohormones plays a crucial role in regulating plant growth, development, and adaptive responses. Red, blue, and far-red light influence the synthesis and activity of auxins, gibberellins, cytokinins, abscisic acid, and ethylene, mediating processes such as elongation, leaf expansion, and stress responses. Understanding these mechanisms enables optimized crop management in controlled environments like greenhouses and vertical farms, promoting higher yield and quality. Integrating light-hormone knowledge into practical agriculture offers sustainable solutions for enhancing productivity, especially under limited natural light conditions, while maintaining plant health and structural integrity.

REFERENCES

[1] «A database of scholarly articles on plant physiology».

[2] Y. Meng и others, «Blue Light Regulates Gibberellin Biosynthesis and Plant Height in Rice», *Plant Cell Rep.*, 2013.

[3] A. Lee, «Effects of Red and Blue Light on Plant Growth», *Bot. Res. J.*, сс. 355–362, 2019.

[4] D. Jabborova, «Formation of students' creativity in the classroom of the Russian language», *Linguist. Cult. Rev.*, сс. 467–477, 2021.

[5] H. Chen, «Impact of environmental stress on plant metabolism», *Plant Physiol. Rep.*, сс. 87–94, 2014.

[6] D. Jabborova и X. Zulfiya, «Intertextual elements, their functions in the text (based on

the novel „kys“ by T. Tolstoy)», *Soc. Sci. Innov.*, cc. 90–98, 2023.
[7] W. G. Hopkins и N. P. A. Hüner, *Introduction to Plant Physiology*. Wiley, 2009.
[8] C. Fankhauser и J. Chory, «Light Control of Plant Development», *Curr. Opin. Plant Biol.*, 1997.
[9] J. Smith, «Photosynthesis and Growth Patterns in Tomato Plants», *J. Plant Sci.*, cc. 210–218, 2020.
[10] M. Jones, *Plant Biochemistry*. Oxford University Press, 2018.
[11] D. Roberts, *Plant Cell Biology*. Garland Science, 2016.
[12] L. Taiz и E. Zeiger, *Plant Physiology and Development*. Sinauer Associates, 2015.
[13] P. Brown, *Principles of Plant Nutrition*. Springer, 2017.
[14] D. Jabborova, «Psychological and pedagogical model of development of students' creative abilities in Russian language and literature classes», *Soc. Sci. Innov.*, cc. 84–89, 2023.
[15] L. Wang, «The role of phytochromes in plant photomorphogenesis», *Annu. Rev. Plant Biol.*, cc. 21–44, 2015.